FSA Algebra I End-of-Course Review Packet Algebra and Modeling

Table of Contents

MAFS.912.A-APR.1.1 EOC Practice	3
MAFS.912.A-CED.1.1 EOC Practice	
MAFS.912.A-REI.2.3 EOC Practice	7
MAFS.912.A-CED.1.4 EOC Practice	
MAFS.912.A-CED.1.2 EOC Practice	11
MAFS.912.A-REI.3.5 EOC Practice	
MAFS.912.A-REI.3.6 EOC Practic	14
MAFS.912.A-REI.4.12 EOC Practice	
MAFS.912.A-CED.1.3 EOC Practice	19
MAFS.912.A-REI.1.1 EOC Practice	21
MAFS.912.A-REI.4.11 EOC Practice	
MAFS.912.A-REI.4.10 EOC Practice	27
MAFS.912.A-SSE.2.3 EOC Practice	
MAFS.912.A-SSE.1.1 EOC Practice	31
MAFS.912.A-SSE.1.2 EOC Practice	

MAFS.912.A-APR.1.1 EOC Practice

Level 2	Level 3	Level 4	Level 5
adds two polynomials with integral	adds and subtracts polynomials,	completes an informal	explains closure
coefficients, including adding when	including adding or subtracting when	argument on closure; applies	for polynomials
multiplying a constant to one or	one or both polynomials is multiplied	multiple operations (excluding	
both polynomials using the	by a monomial or binomial, with a	division) when simplifying	
distributive property is required	degree no greater than 1	polynomials	

1. What is the product of the following expression?

 $(3x + 6)^2$

A. $6x^2 + 12$


- B. $9x^2 + 36$
- C. $9x^2 + 18x + 36$
- D. $9x^2 + 36x + 36$
- 2. What is the product of the following expression?

$$2x(x^2 + x - 5)$$

- A. $2x^3 + x 5$
- B. $2x^3 + 2x 10$
- C. $2x^3 + 2x^2 5x$
- D. $2x^3 + 2x^2 10x$
- 3. Which is the simplified form of this expression?


 $(2x+3)(x-6) - 2x^2 + 3x + 30$

- E. $4x^2 6x + 12$
- A. $-2x^2 + 6x + 27$
- B. -6x 12
- C. -6x + 12
- 4. In the diagram at the right, the dimensions of the large rectangle are (3x 1) by (3x + 7) units. The dimensions of the cut-out rectangle are x by 2x + 5 units. Which choice expresses the area of the shaded region, in square units?



- A. $x^2 + 23x 7$ B. $x^2 + 13x - 7$ C. $7x^2 + 23x - 7$
- D. $7x^2 + 13x 7$

5. Given $ax^2 + bx + c = 2(1.2x + 0.3)(x - 0.5) + (0.5x^2 + 2.5x - 1.3)$. What are the values of *a*, *b*, and *c*?



- 6. Which expression is equivalent to 2(3g 4) (8g + 3)?
 - A. -2g 1
 - B. -2g 5
 - C. -2g 7
 - D. -2g 11
- 7. Which expression is equivalent to $(2x^2 + 3)(x + 4)$?
 - A. $2x^3 + 12$
 - B. $2x^2 + 11x + 12$
 - C. $2x^3 + 6x^2 + 4x + 12$
 - D. $2x^3 + 8x^2 + 3x + 12$
- 8. Under what operations is the system of polynomials NOT closed?
 - A. Addition
 - B. Subtraction
 - C. Multiplication
 - D. Division

MAFS.912.A-CED.1.1 EOC Practice

Level 2	Level 3	Level 4	Level 5
writes or chooses a one-	writes or chooses a simple	writes an exponential equation with	employs the modeling
variable linear equation or	exponential (no horizontal or	a horizontal or vertical translation or	cycle when writing an
inequality in a real-world	vertical translation) or a	a quadratic equation; identifies the	equation
context	simple quadratic equation	meaning of the variables	

 There are 60 students going on a field trip to the chocolate factory. The students are from three different classes. Mrs. Hooper's class has 24 students and Mr. Gomez's class has 18 students. Which of the equalities correctly describes the students and could be used to solve for how many students are from Mr. Anderson's class? (Let A = the number of students in Mr. Anderson's class.)

A. A + 18 = 24

- B. A + A + A = 60
- C. 60 18 = A 24
- D. 24 + 18 + A = 60
- 2. The ages of three friends are consecutively one year apart. Together, their ages total 48 years. Which equation can be used to find the age of each friend (where *a* represents the age of the youngest friend)?
 - A. 3a = 48B. a(a + 1)(a + 2) = 48C. a + (a - 1) + (a - 2) = 48D. a + (a + 1) + (a + 2) = 48
 - a) What are the ages of the friends?
 - A. 16, 17, 18
 - B. 15, 16, 17
 - **C**. 14, 15, 16
 - D. 17, 18, 19
- 3. Student council is renting a tent for \$350 for an upcoming student fair. Each student attending the fair will pay \$0.50. All other attendees will pay \$2.25 each. If 200 students attend the fair, which inequality can be used to determine the number of "other" attendees, *a*, needed to cover the cost of the tent?

A. $(0.50)(200) - 2.25a \ge 350.00$ B. $(0.50)(200) + 2.25a \ge 350.00$ C. $0.50a - (2.25)(200) \ge 350.00$ D. $0.50a + (2.25)(200) \ge 350.00$

- 4. A farmer has a rectangular field that measures 100 feet by 150 feet. He plans to increase the area of the field by 20%. He will do this by increasing the length and width by the same amount, x. Which equation represents the area of the new field?
 - A. (100 + 2x)(150 + x) = 18,000
 - B. 2(100 + x) + 2(150 + x) = 15,000
 - C. (100 + x)(150 + x) = 18,000
 - D. (100 + x)(150 + x) = 15,000

- 5. A heart shaped chocolate box is composed of one square and two half circles. The total number of chocolates in the box is calculated by adding the area of a square given by $4x^2$ and the area of a circle approximated by $3x^2$. The company plans to add a small additional box for a promotional campaign containing one row (2x) of chocolates. If the total combined heart shape and small box contain 69 chocolates, which of these equations could be utilized to solve for the number of chocolates in the small box (2x)?
 - A. $4x^{2} + 3x^{2} + 2x = 69$ B. $4x^{2} - 3x^{2} + 2x = 69$ C. $4x^{2} + 3x^{2} - 2x = 69$ D. $4x^{2} - 3x^{2} - 2x = 69$
- 6. An internet business sells U.S. flags for \$16.95 each, plus \$2.50 shipping per flag. Shipping is free, however, on orders where more than \$100.00 of flags are purchased. Which correctly shows the number of flags f that must be purchased to get free shipping?
 - A. 16.95f = 100
 - B. 16.95*f* > 100
 - C. 19.45*f* > 100
 - D. 16.95f + 2.50 > 100
- 7. A scientist is studying wildlife. She estimates the population of bats in her state to be 270,000. She predicts the population to grow at an average annual rate of 2.9%.
 Using the scientist's prediction, create an equation that models the population of bats, *y*, after *x* years.
- Sandy programmed a website's checkout process with an equation to calculate the amount customers will be charged when they download songs.
 The website offers a discount. If one song is bought at the full price of \$1.29, then each additional song is \$.99. State an equation that represents the cost, *C*, when *s* songs are downloaded.
- 9. Ian is borrowing \$1000 from his parents to buy a notebook computer. He plans to pay them back at the rate of \$60 per month. Ken is borrowing \$600 from his parents to purchase a snowboard. He plans to pay his parents back at the rate of \$20 per month.
 - a) Write an equation that can be used to determine after how many months the boys will owe the same amount.
 - b) Determine algebraically and state in how many months the two boys will owe the same amount. State the amount they will owe at this time.
 - c) Ian claims that he will have his loan paid off 6 months after he and Ken owe the same amount. Determine and state if Ian is correct. Explain your reasoning.

MAFS.912.A-REI.2.3 EOC Practice

Level 2	Level 3	Level 4	Level 5
solves linear equations (with variable on	solves linear equations and	solves linear equations in	solves linear equations
one side and simple benchmark fractions	inequalities in one variable,	one variable, including	and inequalities in one
as the coefficient; may require the use of	where the variable is	equations where one	variable, including
the distributive property and adding like	included on both sides of	coefficient is represented	equations with
terms) and inequalities (with a variable	the equal sign or inequality,	by a letter and requires up	coefficients represented
on one side and positive coefficient that	that require up to three	to three steps to isolate the	by letters that require up
may include a simple benchmark fraction	steps to isolate the variable	variable; solves compound	to four steps to isolate
as the coefficient) in one variable	with rational coefficients	inequalities in one variable	the variable

1. Solve for x: 3(2x - 1) - 10 = 8 + 5x

- A. -7
- В. —З
- C. 19
- D. 21

2. Solve for x: 4(x + 5) = 3(x - 2) - 2(x + 2)

- A. x = -1B. x = -4C. x = -6D. x = -10
- 3. Solve: 3(x + 3) > 4(x 4)
 - A. x > 25B. x < 25C. x > -7D. x < -7
- 4. Solve the following inequality for b, showing all of your work carefully and completely. 4b - 12 - 5b < 9b + 8
- 5. What is the value of x in the equation $\frac{3}{4}x + 2 = \frac{5}{4}x 6$?
 - A. -16
 - B. 16
 - C. -4
 - D. 4

6. Fred solved the equation 8(3x - 7) = -6(x + 7) + 4 as shown.

Given	8(3x-7) = -6(x+7) + 4
Step 1	24x - 56 = -6x - 42 + 4
Step 2	24 x - 56 = -6 x + 46
Step 3	30 x = 102
Step 4	$x = \frac{17}{5}$

Fred made an error between Step 1 and Step 2.

Part A: Explain the error Fred made.

Part B: What is the correct solution to the original equation?

MAFS.912.A-CED.1.4 EOC Practice

Level 2	Level 3	Level 4	Level 5
solves a literal linear equation	solves a literal equation that	solves a literal equation that	solves a literal equation that
in a real-world context for a	requires two procedural steps	requires three procedural	requires four procedural steps
variable whose coefficient is 1		steps	

1. The formula for simple interest plus starting principal, where A = amount, P = principal, r = interest rate per period, and t = time, is given below.

A = P + Prt

Which could be used to find the time, t, if the amount, principal, and interest are known?

A.
$$A - P - Pr = t$$

B. $\frac{A - P}{Pr} = t$
C. $\frac{A - Pr}{P} = t$
D. $\frac{A}{P + rt} = t$

- 2. A line is represented by the equation 3x + 2y = 4. What is another way to represent the same line?
 - A. $y = -\frac{3}{2}x + 2$ B. $y = \frac{3}{2}x + 2$ C. $y = \frac{3}{2}x + 4$ D. $y = -\frac{3}{2}x + 4$
- 3. If k = am + 3mx, the value of m in terms of a, k and x can be expressed as

A.
$$m = \frac{k}{a+3x}$$

B. $m = \frac{k-3mx}{a}$
C. $m = \frac{k-am}{3x}$
D. $m = \frac{k-a}{3x}$

4. A formula is expressed as D = a(2 + kt). Express k in terms of D, a and t.

A.
$$k = \frac{D}{a} - 2t$$

B. $k = D - 2at$
C. $k = \frac{D - 2a}{at}$
D. $k = \frac{D - 2a}{t}$

- 5. Tim was asked to solve the equation for x. His solution is shown below.
 - Start: kx = my mxStep 1: kx + mx = myStep 2: x(k + m) = myStep 3: $x = \frac{my}{k+m}$

In which step did Tim make his first mistake when solving the equation?

- A. Step 1
- B. Step 2
- C. Step 3
- D. Tim did not make a mistake.
- 6. Boyle's Law involves the pressure and volume of gas in a container. It can be represented by the formula $P_1V_1 = P_2V_2$. When the formula is solved for P_2 , the result is
 - A. $P_1V_1V_2$
 - $\mathsf{B.} \quad \frac{V_2}{P_1 V_1}$
 - $\mathsf{C.} \quad \frac{P_1 V_1}{V_2}$
 - D. $\frac{P_1V_2}{V_1}$

MAFS.912.A-CED.1.2 EOC Practice

Level 2	Level 3	Level 4	Level 5
writes or chooses a two-	writes or chooses a system of	writes a system of linear equations or	employs the modeling
variable linear equation for	linear equations or writes a single	writes a single equation that has at	cycle when writing
a real-world context with	equation that has at least three	least three variables; correctly	equations that have
integral coefficients	variables with integral coefficients	identifies the meaning of the variables	two variables

1. Kesha is planning to rent a van for her trip to Mt. Rainier. Two of her friends each rented the same type of van from the same car rental company last week. This is what they told her:

John: "The cost of my rental was \$240. The company charged me a certain amount per day and a certain amount per mile. I had the rental for five days and I drove it 200 miles."


Katie: "The cost of my rental was only \$100. I drove it for 100 miles and had it for two days."

Kesha plans to get the same type of van that John and Katie had from the same car rental company. Kesha estimated her trip would be 250 miles, and she would have the vehicle for four days. Let C = cost, M = miles, and D = daysWhich equation could Kesha use to figure out how much her rental would cost?

A. C = 40.00M + 0.20DB. C = 40.00D + 0.20MC. C = 20.00M + 0.40DD. C = 20.00D + 0.40M

- 2. Eddie's Towing Company charges \$40 to hook a vehicle to the truck and \$1.70 for each mile the vehicle is towed. Which equation best represents the relationship between the number of miles towed, m, and the total charges, c?
 - A. c = 40 + 1.70B. c = 40 + 1.70mC. c = 40m + 1.70D. c = 40m + 1.70
- The local deli charges a fee for delivery. On Monday, they delivered two dozen bagels to an office at a total cost of \$8. On Tuesday, three dozen bagels were delivered at a total cost of \$11. Which system of equations could be used to find the cost of a dozen bagels, b, if the delivery fee is f?
 - A. b + 2f = 8 b + 3f = 11B. 2b + f = 8 b + 3f = 11C. b + 2f = 8 3b + f = 11D. 2b + f = 83b + f = 11

- 4. Max purchased a box of green tea mints. The nutrition label on the box stated that a serving of three mints contains a total of 10 Calories.
 - a) On the axes below, graph the function, C, where C (x) represents the number of Calories in x mints.



- b) Write an equation that represents C(x).
- c) A full box of mints contains 180 Calories. Use the equation to determine the total number of mints in the box.
- A shipping company charges \$1.20 times the sum, s, of the length, width, and height of a package to be shipped. All dimensions are measured in inches. The company also charges \$3.00 for processing the package to be shipped. On the line below, write an equation that the shipping company can use for determining the cost, *C*, for shipping any package.

Equation: ______

6. A construction company spends w weeks extending an existing road. The existing road is 5 miles long. Each week the company completes 0.2 miles of the extension. Which equation models the total length (L) of the road over time?

A. L = 0.22 + 5B. L = 0.22 - 5C. w = 0.22 + 5

D. w = 0.22 - 5

MAFS.912.A-REI.3.5 EOC Practice

Level 2	Level 3	Level 4	Level 5
identifies an equivalent system of	identifies an equivalent system that	identifies systems that	justifies why multiple
two equations in two variables that	has a sum of the original as one of	have the same	equivalent systems
has a multiple of one of the	the equations and a multiple of the	solutions	would have the same
equations of the original system	other		solution

 The Smith Family Reunion and the Jones Family Reunion both include a visit to a family friendly amusement park in Florida. The Smith family pays \$ 882.00 for passes for 10 adults and 18 children. The Jones family pays \$ 951.00 for passes for 11 adults and 19 children. Which equation below can be used to solve for the price of the adult and child admissions?

A. 882 + 951 = (10A + 11A) + (18C + 19C)B. 882 - 951 = (10A - 11A) + (18C - 19C)C. 882 = 10A - 18C; 951 = 11A - 19C

- -2002 1011 100, 551 1111 100
- D. 882 = 10A + 18C; 951 = 11A + 19C
- 2. Which system of equations has the same solution as the system below?

$$2x + 2y = 16$$
$$3x - y = 4$$

- A. 2x + 2y = 16
6x 2y = 4B. x + y = 16
3x y = 4C. 2x + 2y = 16
6x 2y = 8D. 6x + 6y = 48
6x + 2y = 8
- 3. Without solving the systems, explain why the following systems must have the same solution.

System (a): 4x - 5y = 133x + 6y = 11System (b): 8x - 10y = 26x - 11y = 2

4. Which pair of equations could not be used to solve the following equations for x and y?

$$4x + 2y = 22$$
$$-2x + 2y = -8$$

A. 4x + 2y = 22
2x - 2y = 8B. 12x + 6y = 66
6x - 6y = 24C. 4x + 2y = 22
-4x + 4y = -16D. 8x + 4y = 44
-8x + 8y = -8

2016-2017 Algebra and Modeling – Student Packet

MAFS.912.A-REI.3.6 EOC Practice

Level 2	Level 3	Level 4	Level 5
solves a system of linear equations	explains whether a system of equations	solves a system of	[intentionally
approximately when given a graph of the	has one, infinitely many, or no solutions;	equations with rational	left blank]
system; solves a system of equations using	solves a system of equations by graphing	coefficients by	
elimination in the form of ax + by = c and	or substitution (manipulation of equations	graphing, substitution,	
dx + ey = f with integral coefficients,	may be required) or elimination in the	or elimination;	
where only one equation requires	form of ax + by = c and dx + ey = f, where	interprets solutions in a	
multiplication; solves a simple system of	multiplication is required for both	real-world context	
equations that require substitution	equations		

- 1. Sandy has a total of 35 coins in her money jar. If Sandy's jar contains only nickels and dimes and the value of all the coins is \$2.50, how many nickels does Sandy have?
 - A. 5
 - B. 15
 - C. 20
 - D. 30
- 2. The enrollment at High School R has been increasing by 20 students per year. Currently High School R has 200 students attending. High School T currently has 400 students, but its enrollment is decreasing in size by an average of 30 students per year. If the two schools continue their current enrollment trends over the next few years, how many years will it take the schools to have the same enrollment?
 - A. 4 years
 - B. 5 years
 - C. 10 years
 - D. 20 years
- 3. What is the solution for the system of equations?

$$y = 2x - 3$$
$$4x - 3y = 31$$

- A. (-11, -25)
 B. (-11, -19)
 C. (11, 19)
- D. (14,25)
- 4. What is the *y*-coordinate in the solution for the system of linear equations below?

$$-3x + 2y = 6$$

$$4x - y = 2$$


A. -6
B. 1
C. 2
D. 6

5. In attempting to solve the system of equations y = 3x - 2 and 6x - 2y = 4, John graphed the two equations on his graphing calculator. Because he saw only one line, John wrote that the answer to the system is the empty set. Is he correct? Explain your answer.

MAFS.912.A-REI.4.12 EOC Practice


Level 2	Level 3	Level 4	Level 5
identifies a solution region	graphs solutions of the system of two linear	verifies ordered pairs	justifies why an
when the graph of a linear	inequalities and identifies the solution set as a	as being a part of the	ordered pair is a part
inequality is given	region of the coordinate plane that satisfies both	solution set of a	of a solution set
	inequalities; if the form is written in ax + by	system of inequalities	
	< c format, then a, b, and c should be integers		

1. Which system of inequalities describes the graph?




- A. y < 2x 3 $y \ge -\frac{1}{3}x + 2$
- B. $y \le 2x 3$ $y > -\frac{1}{3}x + 2$
- $\begin{array}{ll} \text{C.} & y > 2x 3 \\ & y \leq -\frac{1}{3}x + 2 \end{array}$
- D. $y \ge 2x 3$ $y < -\frac{1}{3}x + 2$
- 2. Which quadrant will be completely shaded by the graph of the inequality y < 3x?
 - A. Quadrant I
 - B. Quadrant II
 - C. Quadrant III
 - D. Quadrant IV


3. Which is a graph of the solution set of the inequality $3x - 4y \le 24$



4. Which graph best represents the solution to this system of inequalities? $\begin{cases} 2x \ge y - 1 \\ 2x - 5y \le 10 \end{cases}$



Β.



5. Without graphing, which point is a solution to the system below?

$$2y < -12x + 4$$
$$y < -6x + 4$$

- A. $(1, \frac{1}{2})$
- B. (0,6)
- C. $\left(-\frac{1}{2}, 5\right)$
- D. (-3,2)


MAFS.912.A-CED.1.3 EOC Practice

Level 2	Level 3	Level 4	Level 5
identifies constraints that are	identifies variables; writes	models constraints using a combination	employs the
constant values or simple	constraints as a system of	of linear equations/inequalities;	modeling cycle when
linear equations/inequalities	linear inequalities or linear	interprets solutions as viable or	writing constraints
in a real-world context	equations	nonviable based on the context	

- On the day of the field trip, each teacher must call the parents of any student who has not returned a permission slip. All of Mr. Gomez's students returned their permission slips, so he did not have to make any calls. Mrs. Hooper and Mr. Anderson had to call a total of eight parents. Mrs. Hooper needed to call two more students than Mr. Anderson. Which set of equations correctly describes the phone calls made? (Let H = Mrs. Hooper's calls and A = Mr. Anderson's calls.)
 - A. H + A = 8; H = A + 2B. H + A = 8; A = H + 2C. H + A = 2; H = A + 8D. H + A = 2; A = H + 8
- 2. In a basketball game, Marlene made 16 fields goals. Each of the field goals were worth either 2 points or 3 points, and Marlene scored a total of 39 points from field goals.

Part A

Let x represent the number of two-point field goals and y represent the number of three-point field goals. Which equations can be used as a system to model the situation? Select **ALL** that apply.



Part B

How many three-point field goals did Marlene make in the game? Enter your answer in the box.

- 3. Justin plans to spend \$20 on sports cards. Regular cards cost \$3.50 per pack and foil cards cost \$4.50 per pack. Which inequality shows the relationship between the number of packs of regular cards (*r*) and the number of packs of foil cards (*f*) Justin can afford to buy?
 - A. $3.5f + 4.5r \le 20$
 - B. $3.5r + 4.5f \le 20$
 - C. $3.5f + 4.5r \ge 20$
 - D. $3.5r + 4.5f \ge 20$
- 4. The amount of profit, p, you earn by selling knives, k, can be determined by: p = 200k 500
 - a) Determine the constraints on profit and the constraints on the number of knives sold.
 - b) What happens to your profit as you sell more knives?
 - c) Is it possible to make a \$14,000 profit? Explain.
- 5. Two friends went to a restaurant and ordered one plain pizza and two sodas. Their bill totaled \$15.95. Later that day, five friends went to the same restaurant. They ordered three plain pizzas and each person had one soda. Their bill totaled \$45.90.

Write and solve a system of equations to determine the price of one plain pizza.

MAFS.912.A-REI.1.1 EOC Practice

Level 2	Level 3	Level 4	Level 5
chooses the correct	chooses the correct	explains and justifies the steps	explains and justifies the steps
justifications for the steps	justifications for the steps in an	in an equation of the form	in an equation of the form a(bx
in a two-step equation,	equation of the form a(bx +c) =	a(bx + c) = d or ax + b = cx + d,	+c) = d(ex +f) , where a, b, c, d,
ax + b = c	d or ax + b = cx + d, where a, b,	where a, b, c, and d are rational	e, and f are rational numbers
	c, and d are integers	numbers	

- 1. State the missing steps and reasons to this solution of 3(x + 4) = 18.
- 2. John's solution to an equation is shown below.
 - Given: $x^2 + 5x + 6 = 0$ Step 1: (x+2)(x+3) = 0Step 2: x+2=0 or x+3=0Step 3: x=-2 or x=-3

Which property of real numbers did John use for Step 2?

- A. multiplication property of equality
- B. zero product property of multiplication
- C. commutative property of multiplication
- D. distributive property of multiplication over addition

3. Which equations illustrate the zero property of multiplication? Select ALL that apply.

```
\boxed{\begin{array}{c}1\\3} \cdot 3 - 3 = 4\\\\\hline\\1\\2 + 2 - 2 = \frac{1}{2}\\\\\hline\\0 \cdot \frac{1}{9} \cdot 9 = 0\\\\\hline\\x - 5 + 5 = x\\\\\hline\\0 \cdot (9 + 3) = 0\end{array}}
```

For questions 4 and 5, use the solution to the equation 3(x - 9) = 12 below.

- Start: 3(x 9) = 12Step 1: 3x - 27 = 12Step 2: 3x - 27 + 27 = 12 + 27Step 3: 3x = 39Step 4: x = 13
- 4. In Step 1, the multiplication property of equality was applied.
 - True
 - False
- 5. In Step 3, the addition property of equality was applied.
 - True
 - False
- 6. Use the steps in the table to answer the question.

initial equation	$3(x+2)^2 + 6x - x$	=	25x + 7x
step 1	$3(x+2)^2 + 5x$	=	32 <i>x</i>
step 2	$3(x+2)^2$	=	27 <i>x</i>
step 3	$(x+2)^2$	=	9 <i>x</i>
step 4	$x^2 + 4x + 4$	=	9 <i>x</i>
step 5	$x^2 - 5x + 4$	=	0

The table shows the first 5 steps used to solve an equation.

Which statement is an incorrect explanation of one step in the process?

- A. From step 4, apply the subtraction property of equality to $x^2 + 4x + 4$ and 9x to get $x^2 55 + 4 = 0$.
- B. From step 3, apply the distributive property to $(x + 2)^2$ to get $x^2 + 4x + 4$ in step 4.
- C. From step 2, apply the distributive property to $3(x + 2)^2$ and 27 to get $(x + 2)^2 = 9x$ in step 3.
- D. From step 1, apply the subtraction property of equality to 5x and 32x to get $3(x + 2)^2 = 27x$ in step 2.

2016-2017 Algebra and Modeling – Student Packet

MAFS.912.A-REI.2.4 EOC Practice

Level 2	Level 3	Level 4	Level 5
solves quadratic equations of the form $x^2 + c = d$, where c and d are rational numbers by simple inspection or by taking square roots	solves quadratic equations of the form $x^2 + bx + c = d$, where b , c , and d are integers by completing the square, factoring, or using the quadratic formula; validates why taking the square root of both sides when solving a quadratic will yield two solutions	solves quadratic equations of the form $x^2 + bx + c = d$, where a, b, c , and d are integers and b/a is an even integer; recognizes that a quadratic can yield nonreal solutions and that the quadratic formula is used to find complex solutions; completes steps in the derivation of the quadratic formula	determines if a quadratic will yield complex solutions; derives the quadratic formula

- 1. What is the solution set of the equation (x 2)(x a) = 0?
 - A. -2 and a
 - B. -2 and -a
 - C. 2 and *a*
 - D. 2 and a
- 2. Janice is asked to solve $0 = 64x^2 + 16x 3$. She begins the problem by writing the following steps: Line 1 $0 = 64x^2 + 16x - 3$
 - Line 2 $0 = B^2 + 2B 3$
 - Line 3 0 = (B+3)(B-1)


Use Janice's procedure to solve the equation for x. Explain the method Janice used to solve the quadratic equation.

- 3. Which value of x is a solution to the equation $13 36x^2 = -12$?
 - A. $\frac{36}{25}$ B. $\frac{25}{36}$ C. $-\frac{6}{5}$ D. $-\frac{5}{6}$
- 4. The method of completing the square was used to solve the equation $2x^2 12x + 6 = 0$. Which equation is a correct step when using this method?
 - A. $(x 3)^2 = 6$
 - B. $(x 3)^2 = -6$
 - C. $(x 3)^2 = 3$
 - D. $(x 3)^2 = -3$

5. An equation is shown.

$$2x^2 - 5x - 3 = 0$$

What values of x make the equation true?



6. Shannon and Jermaine are solving quadratic equations. This table shows their work.

Steps	Shannon	Jermaine
initial equation	$x^2 - 6x + 5 = 12$	$x^2 + 2x - 29 = 2x + 7$
step 1	$x^2 - 6x = 7$	$x^2 + 2x - 36 = 2x$
step 2	$x^2 - 6x + 9 = 16$	$x^2 - 36 = 0$
step 3	$(x-3)^2 = 16$	(x-18)(x+18) = 0
step 4	$x - 3 = \pm 4$	x - 18 = 0 or $x + 18 = 0$
step 5	$x = \pm 7$	x = 18 or x = -18

Both Shannon and Jermaine have errors in their work. Write a clear explanation of each student's error. Provide the correct solutions for both equations.

Shannon	
Correct solution(s):	


Explanation of error:


Explanation of error

MAFS.912.A-REI.4.11 EOC Practice


Level 2	Level 3	Level 4	Level 5
determines an integral solution for	determines a	completes an explanation	explains how to find an approximate
f(x) = g(x) given a graph or a table of	solution to the	on how to find an	solution to the nearest tenth for f(x)
a linear, quadratic, or exponential	nearest tenth for f(x)	approximate solution to the	= g(x) given a graph or a table and
function, in a mathematical or real-	= g(x) given a graph	nearest tenth for f(x) = g(x)	justifies why the intersection of two
world context	or a table	given a graph or a table	functions is a solution to f(x) = g(x)

1. The system 5y = 8x + 8 and 7y = -8x + 16 is graphed as shown. Which choice is the point of intersection?






2. At which point do the two equations 3x + 5 = y + 4x and $y = x^2$ intersect?



- A. (1.8, 3.2)
- B. (-2.8, 7.8)
- C. (0,5)
- D. *Both* (*A*) *and* (*B*)

3. Use the graph below:



If $f(x_1) = g(x_1)$ and $g(x_2) = h(x_2)$, what is $f(x_1) + g(x_2)$?

A. -3

- B. 0
- C. 3
- D. 4

For questions 4and 5, use the table below.

x	-4	-3	-2	-1	0	1
f(x)	-23	-10	-3	-2	-7	-18
<i>g</i> (<i>x</i>)	-13	-11.5	-10	-8.5	-7	-5.58


4. f(x) = g(x) at (0, -7)

- □ True
- □ False
- 5. f(x) = g(x) somewhere on the interval -3 < x < -2.
- True
- □ False

MAFS.912.A-REI.4.10 EOC Practice

Level 2	Level 3	Level 4	Level 5
distinguishes between coordinates	distinguishes between coordinates	recognizes that a graph is	justifies that a graph is
that are solutions to linear	that are solutions to equations in	the set of all the solutions	the set of all the solutions
equations in two variables and	two variables (quadratic or	of a given equation	of an equation
those that are not	exponential) and those that are not		

1. The ordered pairs (20, -29.5), (21, -31), and (22, -32.5) are points on the graph of a linear equation. Which of the following graphs show all of the ordered pairs in the solution set of this linear equation?




- 2. Dr. Math thinks he knows more than you about what is true and false world just because he's a doctor. He says that the equation y = 17x + 1 also includes the point (1, 8). Is Dr. Math right or wrong?
 - A. He's right
 - B. He's wrong
 - C. We need more information before we can say if he's right or wrong
 - D. None of the above

- 3. You talk on the phone y minutes on day x of every month according to the equation y = 2x + 1. The cell phone company claims you talked 12 minutes on the phone on the fourth day of the month. Are they right?
 - A. Yes, you did talk on the phone for 12 minutes on the fourth of the month
 - B. No, you talked on the phone for 7 minutes on the fourth of the month
 - C. No, you talked on the phone for 9 minutes on the fourth of the month
 - D. No, you talked on the phone for 15 minutes on the fourth of the month
- 4. The speed of a snowboarder from uphill to downhill can be modeled using the equation $y = x^2 + 1$ where x is in minutes. The snowboarder's speed at time 0 is 1 and is 2 at time 1. The snowboarder claims that this proves his speed increases linearly. Is he right?
 - A. Yes, because two points are needed to define a line
 - B. No, because the equation is not linear
 - C. No, because the two points have positive values only
 - D. No, because it does not cross the x-axis
- 5. Which point is NOT on the graph represented by $y = -x^2 2x + 8$?
 - A. (-4, 0)
 - B. (-1,9)
 - C. (2,0)
 - D. (4,0)
- 6. An equation is shown.

$$y = \frac{1}{2}x + \frac{3}{4}$$

Select All of the points that are solution to the equation above.



MAFS.912.A-SSE.2.3 EOC Practice

Level 2	Level 3	Level 4	Level 5
uses properties of	factors the difference of two	factors the difference of two squares with a	explains the
exponents (one	squares with a degree of 2 and	common integral factor, trinomials with a	differences
operation) and	trinomials with a degree of 2	common integral factor and a leading coefficient	between equivalent
identifies the new	and explains the properties of	having more than four factors and explains the	forms and why an
base of an	the zeros; completes the square	properties of the zeros; completes the square	equivalent form
exponential	when the leading coefficient is 1	when the leading coefficient is greater than 1	would provide the
function; explains	and explains the properties of	and explains the properties of the maximum or	required property
the properties of	the maximum or minimum; uses	minimum; transforms exponential functions that	
the a in $y = ab_x$ in a	the properties of exponents and	have more than one operation and explains the	
real- world context	names the new rate	properties of expression	

- 1. The director of a play must decide how much to charge per ticket. If tickets cost c dollars each, a total of (755c) people will attend the play. Which ticket price will generate the most income?
 - A. \$1.00
 - B. \$7.50
 - C. \$15.00
 - D. \$20.50
- 2. Which of these shows the following expression factored completely?

 $6x^2 + 15x - 36$

- A. (2x 3)(x + 4)
- B. (6x + 9(x 4))
- C. 3(2x-3)(x+4)
- D. 3(2x+3)(x-4)
- 3. If $f(x) = 2x^2 8x + 9$, which statement regarding the vertex form of f(x) is true?
 - A. In vertex form, $f(x) = 2(x 2)^2 + 1$ and therefore has a minimum value of 1.
 - B. In vertex form, $f(x) = 2(x 2)^2 + 1$ and therefore has a minimum value of 2.
 - C. In vertex form, $f(x) = 2(x 2)^2 + 4.5$ and therefore has a minimum value of 4.5.
 - D. In vertex form, $f(x) = 2(x 2)^2 + 4.5$ and therefore has a minimum value of 2.
- 4. Which expression is equivalent to $x^4 12x^2 + 36$?
 - A. $(x^2 6)(x^2 6)$
 - B. $(6-x^2)(6 + x^2)$
 - C. $(x^2 + 6)(x^2 + 6)$
 - D. $(x^2 + 6)(x^2 6)$

- 5. What number should be added to both sides of the equation to complete the square in $x^2 + 8x = 17$?
 - A. 4
 - B. 16
 - C. 29
 - D. 49
- 6. If (x 7) is a factor of $2x^2 11x + k$, what is the value of k?
 - A. -21
 - B. -7
 - C. 7
 - D. 28
- 7. In the equation $y = (x 2)^2$, the minimum value occurs when x is
 - A. -2
 - B. 2
 - C. -4
 - D. 4
- 8. A computer application generates a sequence of musical notes using the function $f(n) = 6(16)^n$, where *n* is the number of the note in the sequence and f(n) is the note frequency in hertz. Which function will generate the same note sequence as f(n)?
 - A. $g(n) = 12(2)^{4n}$
 - B. $h(n) = 6(2)^{4n}$
 - C. $p(n) = 12(4)^{2n}$
 - D. $k(n) = 6(8)^{2n}$

MAFS.912.A-SSE.1.1 EOC Practice

Level 2	Level 3	Level 4	Level 5
interprets coefficients or terms of	interprets factors of	interprets more than one	given an interpretation,
exponential and quadratic	exponential and quadratic	part of an expression	chooses the correct part of
expressions in a real-world context	expressions		the expression

1. Combined estimates for Etosha National Park and the Northwestern Population

Year	Base Year	Estimated Number of Elephants	
1998	3	3,218	
2000	5	3,628	
2002	7	3,721	
2004	9	3,571	

The elephant population in northwestern Namibia and Etosha National Park can be predicted by the expression $2,649(1.045)^b$, where b is the number of years since 1995.

What does the value 2,649 represent?

- A. the predicted increase in the number of elephants in the region each year
- B. the predicted number of elephants in the region in 1995
- C. the year when the elephant population is predicted to stop increasing
- D. the percentage the elephant population is predicted to increase each year
- 2. A store manager begins each shift with the same total amount of money. She keeps \$200 in a safe and distributes the rest equally to the 5 cashiers in the store. This situation can be represented by the function

 $y = \frac{(x-200)}{5}$. What does the variable x represent in this situation?

- A. The total amount of money the manager has at the beginning of a shift
- B. The total amount of money the manager has at the end of a shift
- C. The amount of money each cashier has at the beginning of a shift
- D. The amount of money each cashier has at the end of a shift
- 3. A satellite television company charges a one-time installation fee and a monthly service charge. The total cost is modeled by the function y = 40 + 90x. Which statement represents the meaning of each part of the function?
 - A. *y* is the total cost, *x* is the number of months of service, \$90 is the installation fee, and \$40 is the service charge per month.
 - B. y is the total cost, x is the number of months of service, \$40 is the installation fee, and \$90 is the service charge per month.
 - C. x is the total cost, y is the number of months of service, \$40 is the installation fee, and \$90 is the service charge per month.
 - D. x is the total cost, y is the number of months of service, \$90 is the installation fee, and \$40 is the service charge per month.

- 4. A ball was thrown upward into the air. The height, in feet, of the ball above the ground t seconds after being thrown can be determined by the expression $-16t^2 + 40t + 3$. What is the meaning of the 3 in the expression? Select the correct answer.
 - A. The ball takes 3 seconds to reach its maximum height.
 - B. The ball takes 3 seconds to reach the ground.
 - C. The ball was thrown from a height of 3 feet.
 - D. The ball reaches a maximum height of 3 feet.
- 5. Is the equation $A = 21000(1 0.12)^t$ a model of exponential growth or exponential decay, and what is the rate (percent) of change per time period?
 - A. exponential growth and 12%
 - B. exponential growth and 88%
 - C. exponential decay and 12%
 - D. exponential decay and 88%
- 6. A car leaves Albany, NY, and travels west toward Buffalo, NY. The equation D = 280 59t can be used to represent the distance, D, from Buffalo after t hours. In this equation, the 59 represents the
 - A. car's distance from Albany
 - B. speed of the car
 - C. distance between Buffalo and Albany
 - D. number of hours driving
- 7. Juan buys peaches and grapefruit at the store. He writes the equations shown to model the relationship between the number of pounds of peaches, *p*, and the number of pounds of grapefruit, *g*, that he buys.

$$p + g = 2.5$$

 $1.58p + 1.09g = 3.46$

What is the total number of pounds of peaches and grapefruit that Juan buys?



pounds

8. Omar deposited d dollars into a savings account y years ago. Now he is going to use a portion of the money in his savings account to buy a bicycle. This expression can be used to find the percentage of the money in the savings account that Omar will use for the bicycle.

$$\frac{342}{d(1.03)^y} \times 100$$

What is the meaning of the denominator in the expression?

- A. the amount Omar will pay for the bicycle
- B. the amount in Omar's savings account now
- C. the yearly interest rate for the savings account
- D. the amount originally deposited in the savings account

MAFS.912.A-SSE.1.2 EOC Practice

Level 2	Level 3	Level 4	Level 5
works with expressions with	factors the difference of two	factors the difference of two	factors the difference of two
only monomial factors and	squares with a degree of 2,	squares with a common	squares with a degree of 4
chooses the correct	trinomials with a degree of 2	integral factor, trinomials with	with or without a common
equivalent forms of a	whose leading coefficient has	a common integral factor and	integral factor, and a
trinomial whose leading	no more than 4 factors	a leading coefficient with more	polynomial with a degree of 3
coefficient is 1		than four factors	and a leading coefficient of 1

- 1. Students were asked to write a trinomial that could not be factored using integers.
 - Pat Wrote:
 $x^2 + 3x 10$

 Sam wrote:
 $x^2 + x 12$

 Mel wrote:
 $x^2 + 2x 1$

 Lee wrote:
 $x^2 + 2x 3$

Which student followed the given directions?

- A. Pat
- B. Sam
- C. Mel
- D. Lee
- 2. Identify ALL the factors of this polynomial when it is factored completely.

 $27x^2 - 153x - 90$

- \square 3 П 9 *x* – 5 \Box *x* + 5 \square 3x - 2 \square 3x + 2П 3x - 15П 9x + 6
- 3. Which expression is equivalent to $16x^2 36$?
 - A. 4(2x-3)(2x-3)
 - B. 4(2x+3)(2x-3)
 - C. (4x-6)(4x-6)
 - D. (4x+6)(4x-6)

- 4. Four expressions are shown below.
 - I $2(2x^2 2x 60)$
 - II $4(x^2 x 30)$
 - III 2(x+6)(x-5)
 - IV 4x(x-1) 120

The expression $4x^2 - 4x - 120$ is equivalent to

- A. I and II, only
- B. II and IV, only
- C. I, II, and IV
- D. II, III, and IV
- 5. Which of these shows the following expression factored completely?

$$6x^2 - 13x + 5$$

- A. (3x 1)(2x + 5)
- B. (3x-5)(2x-1)
- C. (3x-1)(2x-5)
- D. (3x-5)(2x+1)
- 6. Select all the expressions that are equivalent to $9x^4 y^2$.
- $\begin{array}{c|c} (3x^2 y)^2 \\ \hline (3x^2)^2 (y)^2 \\ \hline 9(x^2)^2 (y)^2 \\ \hline (9x^2)^2 (y)^2 \\ \hline (3x^2 + y)(3x^2 y) \end{array}$